

Neutrino oscillations in Kerr-Newman space-time

Jun Ren^{1*} and Cheng-Min Zhang²

1 School of Science, Hebei University of Technology, 300130, Tianjin, China

2 National Astronomical Observatories,

Chinese Academy of Sciences, 100012 Beijing, China

(Dated: February 3, 2010)

The mass neutrino oscillation in Kerr-Newman(K-N) space-time is studied in the plane $\theta = \theta_0$, and the general equations of oscillation phases are given. The effect of the rotation and electric charge on the phase is presented. Then, we consider three special cases: (1) The neutrinos travel along the geodesics with the angular momentum $L = aE$ in the equatorial plane. (2) The neutrinos travel along the geodesics with $L = 0$ in the equatorial plane. (3) The neutrinos travel along the radial geodesics at the direction $\theta = 0$. At last, we calculate the proper oscillation length in the K-N space time. The effect of the gravitational field on the oscillation length is embodied in the gravitational red shift factor. When the neutrino travels out of the gravitational field, the blue shift of the oscillation length takes place. We discussed the variation of the oscillation length influenced by the gravitational field strength, the rotation a^2 and charge Q .

PACS number: 95.30.Sf, 14.60.Pq

Keywords: neutrino oscillation; Kerr-Newman space-time; oscillation length

I. INTRODUCTION

Mass neutrino mixing and oscillations were proposed by Pontecorvo[1], and Mikheyev, Smirnov and Wolfenstein (MSW for short) explored the effect of transformation of one neutrino flavor into another in a medium with varying density[2, 3]. Recently, the consideration of the mass neutrino oscillations has been a hot topic. There have been many

* Email: renjun@hebut.edu.cn

theoretical[4–11] and experimental[12–18] studies about the neutrino oscillations. Then, the neutrino oscillations in the flat space time were extended to the cases in the curved space-time[19–26]. Neutrino oscillation experiments were also considered to test the equivalence principle[27]. Calculating the phase along the geodesic line will produce a factor 2 in the high energy limit, compared with the value along the null line, which often exists in the flat and Schwarzschild space-time[23, 24, 28–31]. This issue of the factor 2 is due to the difference between the time-like and null geodesics. Furthermore, some alternative mechanisms have been proposed to account for the gravitational effect on the neutrino oscillation[32–34]. The inertial effects on neutrino oscillations and neutrino oscillations in non-inertial frames were also called attention[35, 36]. As a further theoretical exploration, neutrino oscillations in space-time with both curvature and torsion[37–39] have been studied.

In recent years, the researches about the neutrino oscillation have been made new progress. A further mechanism to generate pulsar kicks, which was based on the spin flavor conversion of neutrinos propagating in a gravitational field, and the neutrino geometrical optics in gravitational field (in particular in a Lense-Thirring background), have been proposed by Lambiase[40, 41]. Some publications were centered on the theoretical study and experimental measurement of the mixing angle θ_{13} [42–44]. And CP violation in neutrino oscillations were considered by some authors[45–48]. In addition, Cuesta and Lambiase studied the neutrino mass spectrum[49]. Akhmedov, Maltoni and Smirnov presented the neutrino oscillograms for different oscillation channels and discussed the effects of non-vanishing 1-2 mixing[50].

In this paper, we extend the mass neutrino oscillation work from Schwarzschild space-time to Kerr-Newman space-time, since the Kerr-Newman metric is rather important in black hole physics, where a most generally stationary solution with axial symmetry has been existing[51]. For the reason of simplicity, we confine our treatment in two generation neutrinos (electron and muon). We give the general equations of the oscillation phases along the arbitrary null and the time-like geodesics, respectively, in the equal θ plane, $\theta = \theta_0$. The phase along the geodesic will also produce a factor of 2 in the K-N space-time, $\Phi(\text{geod}) = 2\Phi(\text{null})$, in the high energy limit. In our derivation we have not assumed a weak field approximation.

We discuss three spacial cases. Firstly, the oscillation phases along the geodesics with $L = aE$ are considered in the equatorial plane. E is the energy per unit mass of the particle. L and a are the angular momentum per unit mass of the particle and K-N space-time, respectively. The geodesics with $L = aE$ in K-N space-time play the same roles as the radial geodesics in the Schwarzschild and in the Reissner-Nordstrom geometry. In this case, the phases both along the null geodesic and the time-like geodesic are similar in form to the phases along the radial geodesics (null and time-like) in the Schwarzschild space-time. Secondly, we calculate the oscillation phases along the geodesics with $L = 0$ in the equatorial plane. This kind of geodesics is also important in K-N space-time. In the Schwarzschild space-time with non-rotating spherically symmetry, particles with $L = 0$ can propagate along the radial geodesics. But in K-N space-time, because of dragging effect, the coordinate φ must change if a particle with $L = 0$ travels along the geodesics. Thirdly, the phases along the radial geodesics at the direction $\theta = 0$ are given. Only at the poles $\theta = 0$ and $\theta = \pi$, the ergosphere coincides with the event horizon. At the direction $\theta = 0$, the effects of the rotation of the space-time on the oscillation length are found to be more than those in the other directions.

At last, we calculate the proper oscillation length in the K-N space time. The oscillation length is proportional to the local energy (local measurement), $E^{loc} = E/\sqrt{g_{00}}$, of the neutrino, where E is a constant along the geodesic. The decrease in the local energy leads to the decrease in the oscillation length as the neutrino travels out of the gravitational field. So, the blue shift of the oscillation length occurs, which is unlike the case of the gravitational red shift for light signal. In the equatorial plane in K-N space-time, the rotation have no contribution to the oscillation length because g_{00} has nothing to do with the rotating parameter a in this plane. The rotation a^2 of the gravitational field shortens the oscillation length in other equal θ plane, compared with the length in R-N space time. We also give that the length varies according to θ . And charge Q shortens it too, compared with the Kerr space-time case. But, the gravitational field lengthens it, compared with the case in flat space-time.

In this paper, we take the neutrino as a spin-less particle to go along the geodesic because the spin and the curvature coupling has a little contribution to the geodesic derivation[52].

Moreover, the neutrino is a high energy particle, so we do not think the neutrino spin has more contribution to the geodesic.

The paper is organized as the follows. In Sec.2, we briefly review the standard treatment of neutrino oscillation in the flat space-time. In Sec.3, we give the general expressions of the oscillation phases along the null and time-like geodesics in arbitrary equal $\theta = \theta_0$ plane. In Sec.4, we discuss the neutrino phase in three special cases. In Sec.5, we discuss the proper oscillation length in K-N space-time. At last, the conclusion and discussion are given. Throughout the paper, the units $G = c = \hbar = 1$ and $\eta_{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ are used.

II. THE STANDARD TREATMENT OF NEUTRINO OSCILLATION IN FLAT SPACE-TIME

In a standard treatment, the flavor eigenstate $|\nu_\alpha\rangle$ is a superposition of the mass eigenstates $|\nu_k\rangle$, i.e.[21, 22]

$$|\nu_\alpha\rangle = \sum_k U_{\alpha k} \exp[-i\Phi_k] |\nu_k\rangle, \quad (1)$$

where

$$\Phi_k = E_k t - \vec{p}_k \cdot \vec{x}, \quad (k = 1, 2), \quad (2)$$

and the matrix elements $U_{\alpha k}$ comprise the transformation between the flavor and mass bases. E_k and $\vec{p}_k$ are the energy and momentum of the mass eigenstates $|\nu_k\rangle$, and they are different for different mass eigenstates. If the neutrino produced at a space-time point $A(t_A, \vec{x}_A)$ and detected at $B(t_B, \vec{x}_B)$, the expression for the phase in Eq.(2), which is coordinate independent and suitable for application in a curved space-time, is[21, 53]

$$\Phi_k = \int_A^B p_\mu^{(k)} dx^\mu, \quad (3)$$

where

$$p_\mu^{(k)} = m_k g_{\mu\nu} \frac{dx^\nu}{ds}, \quad (4)$$

is the canonical conjugate momentum to the coordinate x^μ and m_k is the rest mass corresponding to mass eigenstate $|\nu_k\rangle$. $g_{\mu\nu}$ and s are metric tensor and an affine parameter, respectively.

The following assumptions are often applied in the standard treatment[4]: (1) The mass eigenstates are taken to be the energy eigenstates, with a common energy; (2) up to $O(m/E)$, there is the approximation $E \gg m$; (3) a massless trajectory is assumed, which means that the neutrino travels along the null trajectory. In the case of two neutrinos mixing $\nu_e - \nu_\mu$, we can write

$$\nu_e = \cos\theta\nu_1 + \sin\theta\nu_2, \nu_\mu = -\sin\theta\nu_1 + \cos\theta\nu_2. \quad (5)$$

Here θ is the vacuum mixing angle. The oscillation probability that the neutrino produced as $|\nu_e\rangle$ is detected as $|\nu_\mu\rangle$ is[54]

$$P(\nu_e \rightarrow \nu_\mu) = |\langle \nu_e | \nu_\mu(x, t) \rangle|^2 = \sin^2(2\theta) \sin^2\left(\frac{\Phi_{kj}}{2}\right), \quad (6)$$

where, $\Phi_{kj} = \Phi_k - \Phi_j$, is the phase shift. From the standard treatment of the neutrino oscillation[21–23], the standard result for the phase is

$$\Phi_k \simeq \frac{m_k^2}{2E_0} |\vec{x}_B - \vec{x}_A|. \quad (7)$$

Here E_0 is the energy for a massless neutrino. So, the phase shift responsible for the oscillation is given by

$$\Phi_{kj} \simeq \frac{\Delta m_{kj}^2}{2E_0} |\vec{x}_B - \vec{x}_A|, \quad (8)$$

where $\Delta m_{kj}^2 = m_k^2 - m_j^2$.

III. NEUTRINO OSCILLATION PHASE ALONG THE NULL AND THE TIME-LIKE GEODESIC IN THE PLANE $\theta = \theta_0$

In this section, we study the neutrino oscillation in equal $\theta = \theta_0$ surface. The line element of K-N space time takes the form

$$ds^2 = g_{00}dt^2 + g_{11}dr^2 + g_{22}d\theta^2 + g_{33}d\varphi^2 + 2g_{03}dtd\varphi. \quad (9)$$

The relevant components of the canonical momentum of the k^{th} massive neutrino in Eq.(4) are

$$\begin{aligned} p_t^{(k)} &= p_0^{(k)} = m_k g_{00} \dot{t} + m_k g_{03} \dot{\varphi}; \\ p_r^{(k)} &= m_k g_{11} \dot{r}; \\ p_\varphi^{(k)} &= m_k g_{33} \dot{\varphi} + m_k g_{03} \dot{t}, \end{aligned} \quad (10)$$

where $\dot{t} = \frac{dt}{ds}$, $\dot{r} = \frac{dr}{ds}$, $\dot{\varphi} = \frac{d\varphi}{ds}$. Because the metric tensor components do not depend on the coordinate t and φ , their canonical momenta $p_t^{(k)}$ and $p_\varphi^{(k)}$ are constant along the trajectory. In fact, the momentum $p_0^{(k)}$ conjugate to t is the asymptotic energy of the neutrino at $r = \infty$. It is stressed that it is the covariant energy p_0 (not p^0) the constant of motion. Otherwise the ambiguous definition of the energy will lead to the confusion in understanding the neutrino oscillation.

The phase along the null geodesic from point A to point B is given by [21, 23, 53]

$$\begin{aligned}\Phi_k^{null} &= \int_A^B p_\mu^{(k)} dx^\mu = \int_A^B (p_0^{(k)} dt + p_\varphi^{(k)} d\varphi + p_r^{(k)} dr) \\ &= \int_A^B (p_0^{(k)} \frac{dt}{dr} + p_\varphi^{(k)} \frac{d\varphi}{dr} + p_r^{(k)}) dr.\end{aligned}\quad (11)$$

We can obtain the following relations which are useful in the calculation

$$\begin{aligned}g^{00} &= -\frac{g_{33}}{\Delta \sin^2 \theta}, g^{33} = -\frac{g_{00}}{\Delta \sin^2 \theta}, \\ g^{03} &= \frac{g_{03}}{\Delta \sin^2 \theta}, g_{03}^2 - g_{00}g_{33} = \Delta \sin^2 \theta,\end{aligned}\quad (12)$$

where $\Delta = r^2 - 2Mr + a^2 + Q^2$. Solving the equation (10) for $\dot{t}$ and $\dot{\varphi}$, we obtain

$$\dot{t} = -\frac{g_{33}E_k + g_{03}L_k}{\Delta \sin^2 \theta_0}, \dot{\varphi} = \frac{g_{03}E_k + g_{00}L_k}{\Delta \sin^2 \theta_0}, \quad (13)$$

where $E_k = \frac{p_0^{(k)}}{m_k}$ and $L_k = -\frac{p_\varphi^{(k)}}{m_k}$ are the energy and angular momentum per unit mass, respectively.

In the standard treatment of the neutrino oscillation, the neutrino is usually supposed to travel along the null[4, 21, 22, 54–56]. Following the standard treatment, we will calculate the phase along the light-ray trajectory from A to B .

The lagrangian appropriate to motions in the plane (for which $\dot{\theta} = 0$ and $\theta = \text{a constant} = \theta_0$) is[57]

$$\mathcal{L} = \frac{1}{2}(g_{00}\dot{t}^2 + 2g_{03}\dot{t}\dot{\varphi} + g_{11}\dot{r} + g_{33}\dot{\varphi}^2). \quad (14)$$

The Hamiltonian is given by

$$\mathcal{H} = E_k \dot{t} - L_k \dot{\varphi} + \frac{p_r^{(k)}}{m_k} \dot{r} - \mathcal{L}. \quad (15)$$

Because of the independence of the Hamiltonian on t , we can deduce that

$$2\mathcal{H} = E_k \dot{t} - L_k \dot{\varphi} + \frac{p_r^{(k)}}{m_k} \dot{r} = \delta_1 = \text{constant}. \quad (16)$$

Without loss generality, we can set, $\delta_1 = 1$ for time-like geodesics, $\delta_1 = 0$ for null geodesics. Substituting (13) into (16) and setting $\delta_1 = 0$ for null geodesics, we have the radial equation

$$g_{11}\dot{r}^2 = \frac{g_{33}E_k^2 + 2g_{03}E_kL_k + g_{00}L_k^2}{\sin^2\theta_0\Delta}. \quad (17)$$

We define a new function

$$V(r) = g_{33}E_k^2 + 2g_{03}E_kL_k + g_{00}L_k^2. \quad (18)$$

The different $V(r)$ determines the phase of the different trajectory. From (17), we get

$$\dot{r} = \frac{\sqrt{-V}}{\rho\sin\theta_0}, \quad (19)$$

where $\rho^2 = r^2 + a^2\cos^2\theta_0$. So, the equations governing t and φ are

$$\frac{dt}{dr} = -\frac{\rho(g_{33}E_k + g_{03}L_k)}{\Delta\sin\theta_0\sqrt{-V}}, \quad \frac{d\varphi}{dr} = \frac{\rho(g_{03}E_k + g_{00}L_k)}{\Delta\sin\theta_0\sqrt{-V}}. \quad (20)$$

The mass-shell condition is[21]

$$m_k^2 = g_{\mu\nu}p^{(k)\mu}p^{(k)\nu} = p_\mu^{(k)}p^{(k)\mu} = p_0^{(k)}p^{(k)0} + p_\varphi^{(k)}p^{(k)\varphi} + p_r^{(k)}p^{(k)r}. \quad (21)$$

Substituting $p^{(k)0} = g^{00}p_0^{(k)} + g^{03}p_\varphi^{(k)}$, $p^{(k)\varphi} = g^{33}p_\varphi^{(k)} + g^{30}p_0^{(k)}$ and (10) into the equation of the mass-shell condition(21), we obtain

$$p^{(k)r} = \frac{m_k\sqrt{-V - \sin^2\theta_0\Delta}}{\rho\sin\theta_0}. \quad (22)$$

In the process of calculation, the relations (12) are used. Applying the relativistic condition $p_0^k \gg m_k$, we have the relation

$$p^{(k)r} \simeq \frac{m_k}{\rho\sin\theta_0}(\sqrt{-V} - \frac{\sin^2\theta_0\Delta}{2\sqrt{-V}}). \quad (23)$$

Adopting (20) and $p^{(k)r}$, the phase along the null geodesics (11) is approximated by

$$\Phi_k^{null} \simeq \int_A^B \frac{m_k\rho\sin\theta_0 dr}{2\sqrt{-V}}. \quad (24)$$

The phase (24) is a general result. The different function $V(r)$ corresponds to the different motion and determines the different phase consequently. If $a = 0$, we can obtain the oscillation phase in the Reissner-Nordstrom space-time; if $Q = 0$, the Kerr space-time case is given. If $a = 0, Q = 0$, the function $V(r)$ reduces to

$$V(r) = -r^2\sin^2\theta_0E_k^2. \quad (25)$$

The phase (24) becomes to

$$\Phi_k^{null} = \int_A^B \frac{m_k^2}{2p_0^{(k)}} dr = \frac{m_k^2}{2p_0^{(k)}} (r_B - r_A). \quad (26)$$

This is just the phase in Schwarzschild space-time[21, 22].

The velocity of an extremely relativistic neutrino is nearly the speed of light. In the standard treatment, the neutrino is supposed to travel along the null line[4, 21, 22, 54–56]. Despite of this, the propagation difference between a massive neutrino and a photon can have important consequences and this tiny derivation becomes important for the understanding of the neutrino oscillation. Therefore, for more general situations, we start to calculate the phase along the time-like geodesic. The factor of 2 will be obtained, when compared the time-like geodesic phase with the null geodesic phase in the high energy limit. The classical orbit is defined to a plane[21, 23], $\theta = \theta_0$, $d\theta = 0$. The phase along the time-like geodesic is[19, 23, 28, 53]

$$\Phi_k^{geod} = \int_A^B p_\mu^{(k)} dx^\mu = \int_A^B (p_0^{(k)} \frac{dt}{dr} + p_\varphi^{(k)} \frac{d\varphi}{dr} + p_r^{(k)}) dr. \quad (27)$$

For time-like geodesic, $\delta_1 = 1$, equation (16) becomes

$$E_k \dot{t} - L_k \dot{\varphi} + \frac{p_r^{(k)}}{m_k} \dot{r} = 1, \quad (28)$$

while the equations for $\dot{t}$ and $\dot{\varphi}$ (13) are the same for time-like geodesics[57]. Substituting $\dot{t}$ and $\dot{\varphi}$, we have

$$\frac{ds}{dr} = \frac{1}{\dot{r}} = \frac{\sqrt{-g_{11}}}{\sqrt{-1 - \frac{V}{\Delta \sin^2 \theta_0}}}. \quad (29)$$

So, we obtain the equations for dt/dr and $d\varphi/dr$ for time-like geodesics

$$\frac{dt}{dr} = -\frac{\sqrt{-g_{11}}}{\Delta \sin^2 \theta_0} \frac{(g_{33} E_k + g_{03} L_k)}{\sqrt{-1 - \frac{V}{\Delta \sin^2 \theta_0}}}, \quad \frac{d\varphi}{dr} = \frac{\sqrt{-g_{11}}}{\Delta \sin^2 \theta_0} \frac{(g_{00} L_k + g_{03} E_k)}{\sqrt{-1 - \frac{V}{\Delta \sin^2 \theta_0}}}. \quad (30)$$

According to mass shell condition, $p_r^{(k)}$ is given by

$$p_r^{(k)} = -m_k \sqrt{-g_{11}} \sqrt{-1 - \frac{V}{\Delta \sin^2 \theta_0}}. \quad (31)$$

Thus, the phase along the time like geodesic is

$$\Phi_k^{geod} = \int_A^B \frac{m_k \sqrt{-g_{11}} dr}{\sqrt{-1 - \frac{V}{\Delta \sin^2 \theta_0}}}. \quad (32)$$

If the high energy limit is taken into account, Eq. (32) reduces to

$$\Phi_k^{geod} \simeq \int_A^B \frac{m_k \rho \sin \theta_0 dr}{\sqrt{-V}} = 2\Phi_k^{null}. \quad (33)$$

It is often noted that the factor 2 of the neutrino phase calculations exists in the flat space-time[29, 30] and in the Schwarzschild space-time[23, 24, 28], which is believed to be the difference between the null geodesic and the time like geodesic. The neutrino phase induced by the null condition, as in the standard treatment, comes from the 4-momentum p^μ defined along the time-like geodesic, and the equation (17) governing $\dot{r}$ to the null geodesic. If the 4-momentum defined along the null geodesic was instead used to compute the null phase, we would obtain zero because of the null condition. When we calculate the phase along the time-like geodesic, $\dot{r}$ in (28) is defined to the time-like geodesic. It is the difference producing the factor 2. It can be proved that the neutrino phase along the null is the half of the value along the time like geodesic in the high energy limit in a general curved space-time(see APPENDIX A in literature[23]).

IV. THREE SPECIAL CASES

A. Oscillation phases along the geodesics with $L_k = aE_k$ in the equatorial plane

It is very important that the geodesic is described in the equatorial plane $\theta = \pi/2$ in the K-N space-time. The geodesics with $L_k = aE_k$ play the same roles as the radial geodesics in the Schwarzschild and in the Reissner-Nordstrom geometry. In this case, for null geodesic $\dot{t}$, $\dot{\varphi}$ and $\dot{r}$ reduce to

$$\dot{t} = \frac{r^2 + a^2}{\Delta} E_k; \dot{\varphi} = \frac{a}{\Delta} E_k; \dot{r} = \pm E_k. \quad (34)$$

These equations in fact define the shear-free null-congruences which we use for constructing a null basis for a description of the K-N space-time in a Newman-Penrose formalism[57]. The function $V(r)$ for null geodesic becomes to,

$$V(r) = -r^2 E_k^2. \quad (35)$$

So, the phase along the null is

$$\Phi_k^{null} \simeq \int_A^B \frac{m_k \rho \sin \theta_0 dr}{2\sqrt{-V}} = \int_A^B \frac{m_k}{2E_k} dr = \frac{m_k^2}{2p_0^{(k)}} (r_B - r_A), \quad (36)$$

which appears the same form as that of the Schwarzschild space-time radial oscillation case.

We now turn to a consideration of the time-like geodesic case. The equations for $\dot{t}, \dot{\varphi}$ are the same as for the null geodesics, while $\dot{r}$ becomes to

$$\dot{r} = \sqrt{E_k^2 + \frac{1}{g_{11}}}. \quad (37)$$

Substituting $L_k = aE_k$ into (32), we obtain the phase along the time-like geodesic

$$\Phi_k^{geod} = \int_A^B \frac{m_k dr}{[(\frac{p_0^{(k)}}{m_k})^2 + \frac{1}{g_{11}}]^{1/2}}. \quad (38)$$

Compared with the phase along the radial time-like geodesic in the Schwarzschild space-time[23],

$$\Phi_k^{geod}(Sch) = \int_A^B \frac{m_k dr}{\sqrt{(\frac{p_0^{(k)}}{m_k})^2 - g_{00}}} = \int_A^B \frac{m_k dr}{\sqrt{(\frac{p_0^{(k)}}{m_k})^2 + \frac{1}{g_{11}}}}, \quad (39)$$

we find that the oscillation phase with $L_k = aE_k$ in K-N space-time has the similar form as the phase along the radial in Schwarzschild space-time. Substituting $g_{11} = -\frac{r^2}{\Delta}$ into equation (38), we have

$$\Phi_k^{geod} = \int_A^B \frac{m_k dr}{\sqrt{b + \frac{2M}{r} - \frac{a^2 + Q^2}{r^2}}}, \quad (40)$$

where $b = (\frac{p_0^{(k)}}{m_k})^2 - 1$. Equation (40) can be integrated directly to give

$$\begin{aligned} \Phi_k^{geod} &= \frac{m_k}{b} \sqrt{br_B^2 + 2Mr_B - a^2 - Q^2} - \frac{m_k}{b} \sqrt{br_A^2 + 2Mr_A - a^2 - Q^2} \\ &\quad - \frac{Mm_k}{b^{3/2}} \ln \frac{br_B + M + \sqrt{b(br_B^2 + 2Mr_B - a^2 - Q^2)}}{br_A + M + \sqrt{b(br_A^2 + 2Mr_A - a^2 - Q^2)}}. \end{aligned} \quad (41)$$

Eq.(41) shows the effects of rotation a^2 on the oscillation phase.

If $a = 0$, we can obtain $\dot{t}, \dot{\varphi}, \dot{r}$ along the radial null-geodesics in the equatorial plane in Reissner-Nordstrom space-time

$$\dot{t} = \frac{r^2}{r^2 - 2Mr + Q^2}, \dot{\varphi} = 0, \dot{r} = \pm E. \quad (42)$$

Therefore, the phases along the radial null and time-like geodesic in Reissner-Nordstrom space-time are given by, respectively

$$\begin{aligned} \Phi_k^{null}(RN) &= \frac{m_k^2}{2p_0^{(k)}}(r_B - r_A), \\ \Phi_k^{geod}(RN) &= \int_A^B \frac{m_k dr}{\sqrt{b + \frac{2M}{r} - \frac{Q^2}{r^2}}}. \end{aligned} \quad (43)$$

Letting $a = 0$ in (41), the integral of equation (43) is given.

B. Oscillation phases along the geodesics with $L = 0$ in the equatorial plane

The geodesics with $L_k = 0$ is another important class of geodesics in K-N space-time. If the coordinate t and φ has a relation $d\varphi/dt = -g_{03}/g_{33}$, the canonical momentum $p_\varphi^{(k)}$ in (10) vanishes. The corresponding $\dot{t}$, $\dot{\varphi}$ and $\dot{r}$ for null geodesic are

$$\dot{t} = -\frac{g_{33}}{\Delta} E_k; \dot{\varphi} = \frac{g_{03}}{\Delta} E_k; \dot{r} = \frac{\sqrt{-g_{33}}}{r} E_k. \quad (44)$$

And $\dot{r}$ for time-like geodesics is

$$\dot{r} = \sqrt{\frac{1 + g_{33} E_k^2 / \Delta}{g_{11}}}. \quad (45)$$

Substituting $L_k = 0$ into (24) and (32), the phases along the null and time-like geodesic are given by, respectively

$$\Phi_k^{null} = \int_A^B \frac{m_k^2}{2p_0^{(k)}} \frac{r dr}{\sqrt{-g_{33}}} = \int_A^B \frac{m_k^2}{2p_0^{(k)}} \sqrt{-g_{11} \widetilde{g_{00}}} dr, \quad (46)$$

$$\Phi_k^{geod} = \int_A^B \frac{\sqrt{-g_{11} \widetilde{g_{00}}} m_k dr}{\sqrt{(\frac{p_0^{(k)}}{m_k})^2 - \widetilde{g_{00}}}}. \quad (47)$$

where $\widetilde{g_{00}} = g_{00} - g_{03}^2/g_{33}$. It is difficult to integrate (46) and (47) directly. We can work out them by expanding as a^2 when a^2 is a small quantity.

C. Oscillation phase along the radial geodesic at $\theta = 0$

Unlike in the Schwarzschild and in the Reissner-Nordstrom space-time, the event horizon does not coincide with the ergosphere where g_{00} vanishes in K-N space-time. This is an important feature which distinguishes the K-N space-time from the others. The ergosphere that is a stationary limit surface coincides with the event horizon only at the poles $\theta = 0$ and $\theta = \pi$. The phase along the null geodesic in the direction $\theta = 0$ can be written as

$$\begin{aligned} \Phi_k^{null} &= \int_A^B \frac{m_k \rho \sin \theta_0 dr}{2\sqrt{-V}} \\ &= \int_A^B \frac{m_k \rho \sin \theta_0 dr}{2\sqrt{r^2 + a^2 + \frac{a^2}{\rho^2}(2Mr - Q^2) \sin^2 \theta_0 \sin \theta_0}}. \end{aligned} \quad (48)$$

Substituting $\theta_0 = 0$, the equation (48) becomes

$$\Phi_k^{null} = \int_A^B \frac{m_k^2}{2p_0^{(k)}} dr = \frac{m_k^2}{2p_0^{(k)}} (r_B - r_A). \quad (49)$$

By similar calculation, the phase along the time-like geodesics at $\theta = 0$ is given by

$$\Phi_k^{geod} = \int_A^B \frac{m_k dr}{(b + \frac{2Mr - Q^2}{r^2 + a^2})^{1/2}}, \quad (50)$$

where $b = (\frac{p_0^{(k)}}{m_k})^2 - 1$.

V. PROPER OSCILLATION LENGTH

The propagation of a neutrino is over its proper distance , but dr in (24) is only a coordinate. The proper distance can be written as[58]

$$dl = \sqrt{(\frac{g_{0\mu}g_{0\nu}}{g_{00}} - g_{\mu\nu})dx^\mu dx^\nu}. \quad (51)$$

In K-N space-time, we have

$$dl = \sqrt{-g_{11}dr^2 + (\frac{g_{03}^2}{g_{00}} - g_{33})d\varphi^2}. \quad (52)$$

Substituting $\frac{d\varphi}{dr}$, we obtain

$$dr = \frac{\sqrt{-g_{00}V}}{\sqrt{\Delta}E_k \sin\theta_0} dl. \quad (53)$$

In order to discuss conveniently, we adopt the differential form of (24)

$$d\Phi_k^{null} = \frac{m_k \rho \sin\theta_0 dr}{2\sqrt{-V}}. \quad (54)$$

Substituting (53), we have

$$d\Phi_k^{null} = \frac{m_k^2}{2p_0^{(k)}} \sqrt{g_{00}} dl. \quad (55)$$

It is assumed that the mass eigenstates are taken to be the energy eigenstates, with a common energy in the standard treatment. The equal energy assumption is considered to be correct by some authors[29, 31, 59] and studied carefully in papers[24, 60, 61]. In addition, it is adopted widely in many literatures, for example[21–23, 62]. p_0 will represent the common energy of different mass eigenstates. In fact, the condition of equal momentum

is also adopted to study the neutrino oscillation. In the flat space-time, both conditions (the equal energy and the equal momentum) present practically the same neutrino oscillation results[24]. There are conditions of time translation invariance and space translation invariance in the flat space-time. So, energy conservation and momentum conservation of a free particle are right. In the curved (stationary) space-time, the energy of a particle is conserved along the geodesic due to the existence of a time-like killing vector field. However, the canonical conjugate momentum to r , p_r is not conserved because $(\frac{\partial}{\partial r})^a$ is not killing in the curved (stationary) space-time. Consequently, it is very difficult to study neutrino oscillation if the condition of equal momentum is adopted in curved space-time. In this section, our discussion is on the base of the results in the standard treatment which the phase is calculated along the null. Then, the phase shift which determines the oscillation is

$$d\Phi_{kj}^{null} = d\Phi_k^{null} - d\Phi_j^{null} = \frac{\Delta m_k^2}{2p_0} \sqrt{g_{00}} dl, \quad (56)$$

where $\Delta m_{kj}^2 = m_k^2 - m_j^2$. The equation (56) can be rewritten as

$$\frac{dl}{d(\frac{\Phi_{kj}^{null}}{2\pi})} = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{g_{00}}} = \frac{4\pi p_0^{loc}}{\Delta m_{kj}^2}. \quad (57)$$

The term $\frac{4\pi p_0}{\Delta m_{kj}^2 \sqrt{g_{00}}}$ in (57) can be interpreted as oscillation length L_{OSC} (which is defined by the proper distance as the phase shift Φ_{kj}^{null} changing 2π) measured by the observer at rest at a position r , and $p_0^{loc} = p_0/\sqrt{g_{00}}$ is the local energy. As $r \rightarrow \infty$, p_0^{loc} approaches to the energy p_0 measured by the observer at infinity. $\frac{4\pi p_0}{\Delta m_{kj}^2}$ is the oscillation length in the flat space-time. Equation (57) is universal significance in curved space-time. In fact, $\sqrt{g_{00}}$ is the gravitational red shift factor which shows the effect of the gravitational field on the oscillation length. Consider two static observers O (the radial coordinate r) and O' (the radial coordinate r'). The oscillation length measured by O and by O' is, respectively

$$L_{OSC}(r) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{g_{00}(r)}}, L_{OSC}(r') = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{g_{00}(r')}}. \quad (58)$$

We can obtain the relation

$$\frac{L_{OSC}(r')}{L_{OSC}(r)} = \frac{\sqrt{g_{00}(r)}}{\sqrt{g_{00}(r')}}. \quad (59)$$

If $r' > r$, we have $L_{OSC}(r') < L_{OSC}(r)$ and blue shift occurs. Physically, the oscillation length is proportional to the local energy of the neutrino. When the neutrino travels out of the

gravitational field, the local energy decreases. Consequently, the neutrino oscillation length decreases and blue shift takes place. From equation (57), the oscillation length increases in the gravitation field because of $0 < g_{00} < 1$ out of the the infinite red shift surface. The effect of the gravitational blue shift on the oscillation length may have the possible observable effect from experiments. In the Schwarzschild space-time, $g_{00} = 1 - 2M/r$, we have

$$L_{OSC}(Sch) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{1 - 2M/r}}. \quad (60)$$

In order to study the influence of Charge on the neutrino oscillation, we consider the oscillation length in the Reissner-Nordstrom space-time

$$L_{OSC}(RN) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{g_{00}}} = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{1 - \frac{2M}{r} + \frac{Q^2}{r^2}}}. \quad (61)$$

Compared with the case in the Schwarzschild space-time, the oscillation length decreases due to the influence of charge Q .

The metric component g_{00} in the K-N space-time is

$$g_{00} = 1 - \frac{2Mr - Q^2}{\rho^2}, \quad (62)$$

where $\rho^2 = r^2 + a^2 \cos^2 \theta$. In the equatorial plane, there is, $g_{00} = 1 - 2M/r + Q^2/r^2$, which is the same as g_{00} in the Reissner-Nordstrom space-time. Thus, it is concluded that the neutrino oscillation length along the geodesics in the equatorial plane in the K-N space-time is identical to that in the Reissner-Nordstrom space-time and the rotating parameter a^2 does not work in this plane. Therefore, we have to select other plane $\theta = \theta_0 \neq \pi/2$ to highlight the effect of rotation on the oscillation length. In the plane $\theta = \theta_0$, the oscillation length can be written as

$$L_{OSC}(K - N) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{1 - \frac{2Mr - Q^2}{r^2 + a^2 \cos^2 \theta_0}}}. \quad (63)$$

It is obvious that the oscillation length decreases too because of the rotation of the gravitational field compared with that in R-N space-time. Letting $Q = 0$ in (63), the oscillation length in Kerr space-time is given by

$$L_{OSC}(Kerr) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{1 - \frac{2Mr}{r^2 + a^2 \cos^2 \theta_0}}}. \quad (64)$$

Comparing with (63), the charge Q shortens the oscillation length. We can obtain that the oscillation length varies with θ_0 by

$$\frac{d}{d\theta_0} L_{OSC}(K - N) = \frac{4\pi p_0}{\Delta m_{kj}^2} (g_{00})^{3/2} \frac{2Mr - Q^2}{(r^2 + a^2 \cos^2 \theta_0)^2} a^2 \sin \theta_0 \cos \theta_0. \quad (65)$$

In K-N space-time, we conclude that the oscillation length increases with θ within $0 < \theta < \pi/2$, and it becomes maximum in the equatorial plane. Then, it decreases with θ within $\pi/2 < \theta < \pi$. At the direction $\theta = 0$ and $\theta = \pi$, the oscillation length occurs minimum,

$$L_{OSC}(K - N) = \frac{4\pi p_0}{\Delta m_{kj}^2} \frac{1}{\sqrt{1 - \frac{2Mr - Q^2}{r^2 + a^2}}}. \quad (66)$$

In summary, the gravitational field lengthens oscillation length; both the rotation a^2 and the charge Q shorten the oscillation length.

VI. CONCLUSION AND DISCUSSION

In this paper, we have given the phase of mass neutrino propagating along the null and the time like geodesic in the gravitational field of a rotating symmetric and charged object, which is described by Kerr-Newman metric. Most astrophysical bodies in universe have rotation and charge generally. Thus the work about the neutrino oscillation in the K-N space time is important and meaningful for the black hole astrophysics. We work out the general formula of oscillation phase on the equal $\theta = \theta_0$ plane with the generality. The phase along the geodesic is the double of that along the null in the high energy limit, which is the same in the cases in flat and Schwarzschild space-time. By setting $\theta = \pi/2$, the phases in the equatorial plane are given. As $a = 0$ or $Q = 0$, we obtain the phases in the R-N space-time or in the Kerr space-time. Moreover, we study three special cases in K-N space-time: geodesics with $L = aE$; geodesics with $L = 0$; radial geodesics at $\theta = 0$. Among them, the geodesics with $L = aE$ have the same importance as the radial geodesics in the Schwarzschild and in the R-N geometry. The phases obtained are very similar in form to the cases along the radial geodesics in the Schwarzschild and in the R-N space-time.

In Sec.5, the proper oscillation length in the K-N space time is studied in detail. We find that oscillation length in curved space-time is proportional to the local energy, which is

regraded as the neutrino "climbs out of the gravitational potential well". So, the blue shift occurs. Then, the effects of rotation and charge of the space-time on the oscillation length are given. Because of the correction of the gravitation field, the oscillation length increases, compared with the flat space time case. However, both the rotation a^2 and the charge Q shorten the oscillation length. It is noted, the rotation has null contribution to the length in the equatorial plane in K-N space-time, because red shift factor is independence of a^2 in this plane. Finally, we remark that our result exists generality, which can be exploited to study the neutrino oscillation near the rotating compact star, neutron star or black hole.

-
- [1] Pontecorvo B 1957 Mesonium and antimesonium *Zh. Eksp. Theor. Fiz.* **33** 549
 - [2] Mikheyev S P and Smirnov A Y 1986 Resonant amplification of oscillations in matter and solar-neutrino spectroscopy *Il Nuovo Cimento C* **9** 17
 - [3] Wolfenstein L 1978 Neutrino oscillations in matter *Phys. Rev. D* **17** 2369
 - [4] Bilenky S and Petcov S T 1987 Massive neutrinos and neutrino oscillations *Rev. Mod. Phys.* **59** 671
 - [5] Bilenky B, Giunti C and Grimus W 1999 Phenomenology of neutrino oscillations *Progress in Particle and Nuclear Physics* **43** 1 (hep-th/9812360)
 - [6] Nussinov S 1976 Solar neutrinos and neutrino mixing *Phys. Lett. B* **63** 201
 - [7] Kayser B 1981 On the quantum mechanics of neutrino oscillation *Phys. Rev. D* **24** 110
 - [8] Giunti C, Kim C W and Lee U W 1991 When do neutrinos really oscillate? Quantum mechanics of neutrino oscillations *Phys. Rev. D* **44** 3635
 - [9] Rich J 1993 Quantum mechanics of neutrino oscillations *Phys. Rev. D* **48** 4318
 - [10] Kiers K, Nussinov S and Weiss N 1996 Coherence effects in neutrino oscillations *Phys. Rev. D* **53** 537 (hep-ph/9506271)
 - [11] Sarkar U and Mann R B 1988 Neutrino Masses in Superstring Theories with Intermediate Scales *International Journal of Modern Physics A* **3** 2165
 - [12] Fukuda Y et al(Super-K) 1998 Evidence for Oscillation of Atmospheric Neutrinos *Phys. Rev. Lett.* **81** 1562

- [13] Apollonio et al(CHOZZ) M 1999 Limits on neutrino oscillations from the CHOOZ experiment *Phys. Lett. B***466** 415
- [14] Eguchi K et al(KamLAND) 2003 First Results from KamLAND: Evidence for Reactor Antineutrino Disappearance *Phys. Rev. Lett.* **90**, 021802
- [15] Ahn M H et al(K2K) 2003 Indications of Neutrino Oscillation in a 250 km Long-Baseline Experiment *Phys. Rev. Lett.* **90**, 041801; 2006 Measurement of neutrino oscillation by the K2K experiment *Phys. Rev. D***74** 072003
- [16] Ahmed S N et al(SNO) 2004 Measurement of the Total Active 8B Solar Neutrino Flux at the Sudbury Neutrino Observatory with Enhanced Neutral Current Sensitivity *Phys. Rev. Lett.* **92** 181301
- [17] Michael D G et al(MINOS) 2006 Observation of Muon Neutrino Disappearance with the MINOS Detectors in the NuMI Neutrino Beam *Phys. Rev. Lett.* **97** 191801
- [18] Aharmin B et al(SNO) 2008 Independent Measurement of the Total Active 8B Solar Neutrino Flux Using an Array of ^3He Proportional Counters at the Sudbury Neutrino Observatory *Phys. Rev. Lett.* **101** 111301
- [19] Ahluwalia D V and Burgard C 1996 Gravitationally Induced Neutrino-Oscillation Phases *Gen. Relativ. Gravit.* **28** 1161 (arXiv:gr-qc/9603008)
- [20] Piriz D D, Roy M and Wudka J 1996 Neutrino oscillations in strong gravitational fields *Phys. Rev. D***54** 1587 (hep-ph/9604403)
- [21] Fornengo N, Giunti C M, Kim C M and Song J 1997 Gravitational effects on the neutrino oscillation *Phys. Rev. D* **56** 1895 (hep-ph/9611231)
- [22] Cardall C Y and Fuller G M 1997 Neutrino oscillations in curved spacetime: A heuristic treatment *Phys. Rev. D* **55** 7960 (hep-ph/9610494)
- [23] Zhang C M and Beesham A 2001 The General Treatment of the High and Low Energy Particle Interference Phase in a Gravitational Field *Gen. Rel. Grav.* **33** 1011 (gr-qc/0004048)
- [24] Zhang C M and Beesham A 2003 On the Mass Neutrino Phase Along the Geodesic Line and the Null Line in Curved and Flat Spacetime *Int. J. Mod. Phys. D* **12** 727
- [25] Pereira J G and Zhang C M 2000 Some Remarks on the Neutrino Oscillation Phase in a Gravitational field *Gen. Rel. Grav.* **32** 1633 (gr-qc/0002066)

- [26] Huang X J and Wang Y J 2003 Interference Phase of Mass Neutrinos in Kerr Space-Time *Commun. Theor. Phys.* **40** 742; 2004 Mass neutrino oscillations in Reissner Nordstrom space time *Chin. Phys.* **13** 1588; 2006 Mass neutrino oscillations in Robertson Walker space time *Chin. Phys.* **15** 229
- [27] Mann R B and Sarkar U 1996 Test of the Equivalence Principle from Neutrino Oscillation Experiments *Phys. Rev. Lett.* **76** 865 (hep-ph/9505353)
- [28] Bhattacharya T, Habib S and Mottola E 1999 Gravitationally induced neutrino oscillation phases in static spacetimes *Phys. Rev. D* **59** 067301 (1999).
- [29] Lipkin H J 2000 Neutrino oscillations as two-slit experiments in momentum space *Phys. Lett. B* **477** 195
- [30] Lipkin H J 1999 Quantum Mechanics of Neutrino Oscillations - Hand Waving for Pedestrians (hep-ph/9901399)
- [31] Grossman Y and Lipkin H J 1997 Flavor oscillations from a spatially localized source: A simple general treatment *Phys. Rev. D* **55** 2760 (hep-ph/9607201)
- [32] Gasperini M 1988 Testing the principle of equivalence with neutrino oscillations *Phys. Rev. D* **38** 2635
- [33] Mureika J R and Mann R B 1996 Mass or gravitationally-induced neutrino oscillations? a comparison of 8B neutrino flux spectra in a three-generation framework *Phys. Lett. B* **368** 112 (hep-ph/9511220)
- [34] Mureika J R and Mann R B 1996 Three-flavor gravitationally induced neutrino oscillations and the solar neutrino problem *Phys. Rev. D* **54** 2761 (hep-ph/9603335).
- [35] Capozziello S and Lambiase G 2000 Inertial effects on neutrino oscillations *Eur. Phys. J. C* **12** 343 (gr-qc/9910016)
- [36] Lambiase G 2001 Neutrino oscillations in non-inertial frames and the violation of the equivalence principle Neutrino mixing induced by the equivalence principle violation *Europ. J. Phys. C* **19** 553
- [37] Alimohammadi M and Shariati A 1999 Neutrino oscillation in a space-time with torsion *Mod. Phys. Lett. A* **14** 267 (gr-qc/9808066)
- [38] Capozziello S, Iovane G and Lambiase G et al 1999 Fermion helicity flip induced by torsion

- field *Europhys. Lett.* **46** 710 (astro-ph/9904199).
- [39] Capozziello S, Lambiase G and Stornaiolo C 2001 Geometric classification of the torsion tensor in space-time *Annalen Phys.* **10** 713 (gr-qc/0101038)
 - [40] Lambiase G 2005 Pulsar Kicks Induced by Spin Flavor Oscillations of Neutrinos in Gravitational Fields *Mon. Not. Roy. Astron. Soc.* **362** 867-871 (astro-ph/0411242)
 - [41] Lambiase G, Papini G, Punzi R and Scarpetta G 2005 Neutrino optics and oscillations in gravitational fields *Phys. Rev. D* **71** 073011 (gr-qc/0503027)
 - [42] Lychkovskiy O 2006 Constraints on neutrino mixing angle θ_{13} and Supernova neutrino fluxes from the LSD neutrino signal from SN1987A (hep-ph/0604113)
 - [43] R. Adhikari, A. Datta and B. Mukhopadhyaya 2007 Neutrino mass scale and the mixing angle θ_{13} for quasidegenerate Majorana neutrinos *Phys. Rev. D* **76** 073003 (hep-ph/0703318)
 - [44] Eby D A, Frampton P H and Matsuzaki S 2009 Predictions of neutrino mixing angles in a T model *Phys. Lett. B* **671** 386 (hep-ph/0810.4899).
 - [45] Klinkhamer F R 2006 Possible new source of T and CP violation in neutrino oscillations *Phys. Rev. D* **73** 057301
 - [46] Schwetz T 2007 What is the probability that θ_{13} and CP violation will be discovered in future neutrino oscillation experiments? *Phys. Lett. B* **648** 54-59 (hep-ph/0612223).
 - [47] Gava J and Volpe C 2008 Collective neutrino oscillations in matter and CP violation *Phys. Rev. D* **78** 083007 (astro-ph/0807.3418).
 - [48] Altarelli G and Meloni D 2009 CP violation in neutrino oscillations and new physics *Nucl. Phys. B* **809** 158-182
 - [49] Cuesta H J M and Lambiase G 2008 Neutrino Mass Spectrum from Gravitational Waves Generated by Double Neutrino Spin-Flip in Supernovae *Astrophysical Journal* **689** 371
 - [50] Akhmedov E K, Maltoni M and Smirnov A Y 2008 Neutrino oscillograms of the Earth: effects of 1-2 mixing and CP-violation *Journal of High Energy Physics* 0806, 072(hep-ph/0804.1466)
 - [51] Ohanian H C and Ruffini R 1994 *Gravitation and Spacetime* (New York: Norton & Company)
 - [52] Audretsch J 1981 Trajectories and spin motion of massive spin-1/2 particles in gravitational fields *J. Phys. A* **14** 411
 - [53] L. Stodolsky 1979 Matter and light wave interferometry in gravitational fields *Gen. Relativ.*

Gravit. **11** 391

- [54] Boehm F and Vogel P 1992 *Physics of Massive Neutrino* (Cambridge University Press, Cambridge)
- [55] Zuber K 1998 On the physics of massive neutrinos *Phys. Rep.* **305** 295 (hep-ph/9811267)
- [56] Bahcall J N 1994 *Neutrino Astrophysics* (Cambridge University Press, Cambridge) 1983
- [57] Chandrasekhar S 1983 *Mathematical Theory of Black Holes* (Clarendon Press, Oxford; Oxford University Press, New York), 326-350.
- [58] Landau L D and Lifshitz E M 1987 *The Classical Theory of Field* (Forth Revised English Edition, Butterworth Heinemann) 235
- [59] Stodolsky L 1998 When the wavepacket is unnecessary *Phys. Rev. D* **58** 036006
- [60] Giunti C 2001 Energy and Momentum of Oscillating Neutrinos *Mod. Phys. Lett. A* **16** 2363 (hep-ph/0104148)
- [61] De Leo S, Ducati G and Rotelli P 2000 Remarks upon the mass oscillation formulas *Modern Physics Letters A* **15** 2057 (hep-ph/9906460)
- [62] Giunti C and Kim C W 2001 Quantum Mechanics of Neutrino Oscillations *Found. Phys. Lett.* **14**, 213 (hep-ph/0011074)